

FIXED SUBGROUPS IN DIRECT PRODUCTS OF SURFACE GROUPS OF EUCLIDEAN TYPE

JIANCHUN WU, ENRIC VENTURA, AND QIANG ZHANG

ABSTRACT. We give an explicit characterization of which direct products G of surface groups of Euclidean type satisfy that the fixed subgroup of any automorphism (or endomorphism) of G is compressed, and of which it is always inert.

1. INTRODUCTION

For a finitely generated group G , let $\text{rk}(G)$ denote its rank, i.e., the minimal number of the generators of G . We let endomorphisms of G act on the left, i.e., $g \mapsto \phi g$, and denote the monoid of endomorphisms of G by $\text{End}(G)$, and the group of automorphisms by $\text{Aut}(G)$. For an arbitrary family $\emptyset \neq \mathcal{B} \subseteq \text{End}(G)$, the *fixed subgroup* of \mathcal{B} is

$$\text{Fix } \mathcal{B} := \{g \in G \mid \phi g = g, \forall \phi \in \mathcal{B}\} = \bigcap_{\phi \in \mathcal{B}} \text{Fix } \phi \leq G.$$

Definition 1.1. A subgroup $H \leq G$ is said to be

- (i) *inert in G* if $\text{rk}(H \cap K) \leq \text{rk}(K)$ for every (finitely generated) $K \leq G$;
- (ii) *compressed in G* if $\text{rk}(H) \leq \text{rk}(K)$ for every (finitely generated) K with $H \leq K \leq G$;
- (iii) *c -bounded in G* if $\text{rk}(H) \leq c \cdot \text{rk}(G)$; for simplicity, when H is 1-bounded in G , we just say it is *bounded in G* .

In [2], Bestvina–Handel solved the famous Scott’s conjecture that the fixed subgroup of an automorphism of a free group F is bounded in F :

Theorem 1.2 (Bestvina–Handel, [2]). *For any automorphism ϕ of a free group F , $\text{rk}(\text{Fix } \phi) \leq \text{rk}(F)$.*

The notions of inertia and compression were introduced in Dicks–Ventura [3], where the authors proved the following extension of Bestvina–Handel’s result.

Theorem 1.3 (Dicks–Ventura, [3]). *Let F be a finitely generated free group, and let \mathcal{B} be a family of injective endomorphisms of F . Then, $\text{Fix } \mathcal{B}$ is inert in F .*

It is obvious that inertia implies compression, and compression implies boundedness; also, a finite intersection of inert subgroups in G is again inert in G . There are several results in the literature studying these properties for fixed subgroups of automorphisms and endomorphisms of free groups, surface groups, and finite direct products of them. We list some as follows.

1991 *Mathematics Subject Classification.* 20F65, 20F34, 57M07.

Key words and phrases. fixed subgroups, intersections, free groups, surface groups, direct products, inert, compressed.

Theorem 1.4 (Bergman, [1]). *Let F be a finitely generated free group, and let $\mathcal{B} \subseteq \text{End}(F)$. Then, $\text{Fix } \mathcal{B}$ is bounded in F .*

Theorem 1.5 (Martino–Ventura, [7]). *Let F be a finitely generated free group, and let $\mathcal{B} \subseteq \text{End}(F)$. Then, $\text{Fix } \mathcal{B}$ is compressed in F .*

It is still open whether $\text{Fix } \phi$ is inert in F for every $\phi \in \text{End}(F)$, a question asked first in [3, Problem 5], and known as the *inertia conjecture*; a small step into this direction can be found in [13].

For surface groups a lot is known too. A *surface group* is the fundamental group $\pi_1(X)$ of a connected, possibly punctured, surface X . To fix notation, we denote Σ_g the closed orientable surface of genus $g \geq 1$, and

$$S_g = \pi_1(\Sigma_g) = \langle a_1, b_1, \dots, a_g, b_g \mid [a_1, b_1] \cdots [a_g, b_g] \rangle$$

its fundamental group; for the non-orientable case, we denote NS_k the connected sum of $k \geq 1$ projective planes, and

$$NS_k = \pi_1(NS_k) = \langle a_1, a_2, \dots, a_k \mid a_1^2 \cdots a_k^2 \rangle$$

its fundamental group. With one or more punctures, we get the finitely generated free groups F_r . Straightforward calculations show that $NS_2 \simeq \langle a, b \mid bab^{-1}a \rangle$; we shall work with this presentation.

Nielsen, Jaco–Shalen, and Zieschang gave the following results.

Theorem 1.6 (Nielsen, [8, 9]; Jaco–Shalen, [4]). *Let G be a surface group. For any $\phi \in \text{Aut}(G)$, $\text{Fix } \phi$ is bounded in G .*

Theorem 1.7 (Zieschang, [14]). *Let G be a closed surface group with $\chi(G) < 0$. For any non-epimorphic endomorphism $\phi \in \text{End}(G)$, $\text{Fix } \phi$ is $\frac{1}{2}$ -bounded in G .*

Some alternative proofs can also be found in [5]. Some years ago, these two theorems were extended in [10] to families of endomorphisms.

Theorem 1.8 (Wu–Zhang, [10]). *Let G be a closed surface group with $\chi(G) < 0$. For any $\mathcal{B} \subseteq \text{Aut}(G)$, $\text{Fix } \mathcal{B}$ is inert in G .*

And, recently, the analog of Theorem 1.5 has been obtained in [13] for surface groups.

Theorem 1.9 (Zhang–Ventura–Wu, [13]). *Let G be a surface group and $\mathcal{B} \subseteq \text{End}(G)$. Then, $\text{Fix } \mathcal{B}$ is compressed in G .*

To our knowledge, there are only a few results of this type for fundamental groups of 3-manifolds; see [6, 11, 12].

Recently, Zhang–Ventura–Wu started investigating in [13] these same questions but within the family of finite direct products of free and surface groups, i.e., groups of the form $G = G_1 \times \cdots \times G_n$, where each building block G_i is either 1, F_r , S_g , or NS_k . Here, the situation is more complicated and interesting new phenomena show up. In order to study the behaviour of fixed subgroups, it becomes relevant to distinguish between the *Euclidean* building blocks, 1, \mathbb{Z} , \mathbb{Z}^2 , \mathbb{Z}_2 , and NS_2 (corresponding, respectively, to the sphere, two-punctured sphere, torus, projective plane, and Klein bottle), and the *hyperbolic* building blocks, F_r with $r \geq 2$, S_g with $g \geq 3$, and NS_k with $k \geq 3$ (corresponding, respectively, to a r -punctured disc, a

connected sum of g tori, and a connected sum of k projective planes). Geometrically, this is, precisely, distinguishing whether the surface has non-negative or negative Euler characteristic, respectively; and algebraically, it is distinguishing whether the building block has non-trivial or trivial center, respectively (see [13, Lemma 4.2]).

Let G be a finite direct product of surface groups. [13, Theorem 4.8] states that mixing the two types of building blocks is the unique possible way to break boundedness of fixed subgroups of automorphisms of G ; more precisely,

Theorem 1.10 (Zhang–Ventura–Wu, [13]). *Let $G = G_1 \times \cdots \times G_n$, where each G_i is a surface group. Then, $\text{Fix } \phi$ is bounded in G for all $\phi \in \text{Aut}(G)$ if and only if the G_i 's are all Euclidean, or all hyperbolic.*

Then, the authors gave necessary conditions on G for all fixed subgroups of automorphisms of G to be compressed in G , or to be inert in G :

Theorem 1.11 (Zhang–Ventura–Wu, [13]). *Let $G = G_1 \times \cdots \times G_n$, where each G_i is a surface group. If $\text{Fix } \phi$ is compressed in G for every $\phi \in \text{Aut}(G)$, then G must be of one of the following forms:*

- (euc1) $G = \mathbb{Z}^p \times \mathbb{Z}_2^q$ for some $p, q \geq 0$; or
- (euc2) $G = NS_2 \times \mathbb{Z}_2^q$ for some $q \geq 0$; or
- (euc3) $G = NS_2 \times \mathbb{Z}^p \times \mathbb{Z}_2$ for some $p \geq 1$; or
- (euc4) $G = NS_2^\ell \times \mathbb{Z}^p$ for some $\ell \geq 1, p \geq 0$; or
- (hyp1) $G = F_r \times NS_3^\ell$ for some $r \geq 2, \ell \geq 0$; or
- (hyp2) $G = S_g \times NS_3^\ell$ for some $g \geq 2, \ell \geq 0$; or
- (hyp3) $G = NS_k \times NS_3^\ell$ for some $k \geq 3, \ell \geq 0$.

Theorem 1.12 (Zhang–Ventura–Wu, [13]). *Let $G = G_1 \times \cdots \times G_n$, where each G_i is a surface group. If $\text{Fix } \phi$ is inert in G for every $\phi \in \text{Aut}(G)$, then G must be of the form (euc1), or (euc2), or (euc3), or (euc4), or*

- (hyp1') $G = F_r$ for some $r \geq 2$; or
- (hyp2') $G = S_g$ for some $g \geq 2$; or
- (hyp3') $G = NS_k$ for some $k \geq 3$.

In the present paper we complete a full characterization of compression and inertia for the Euclidean case (which happens to be the same for automorphisms and for endomorphisms), by proving the following improvements of Theorems 1.11 and 1.12:

Theorem 1.13. *Let $G = G_1 \times \cdots \times G_n$, where each G_i is a surface group of Euclidean type. Then, the following are equivalent:*

- (a) $\text{Fix } \phi$ is compressed in G for every $\phi \in \text{End}(G)$;
- (b) $\text{Fix } \phi$ is compressed in G for every $\phi \in \text{Aut}(G)$;
- (c) G is of the form (euc1), or (euc2), or (euc4).

Theorem 1.14. *Let $G = G_1 \times \cdots \times G_n$, where each G_i is a surface group of Euclidean type. Then, the following are equivalent:*

- (a) $\text{Fix } \phi$ is inert in G for every $\phi \in \text{End}(G)$;
- (b) $\text{Fix } \phi$ is inert in G for every $\phi \in \text{Aut}(G)$;
- (c) G is of the form (euc1), or (euc2).

As an immediate corollary, together with Theorems 1.12, 1.3, and 1.8, we also get a full characterization of inertia for both types of surfaces, but only concerning automorphisms (the endomorphism case depends on the inertia conjecture, which is still open):

Theorem 1.15. *Let $G = G_1 \times \cdots \times G_n$, where each G_i is a surface group. Then, the following are equivalent:*

- (a) *Fix ϕ is inert in G for every $\phi \in \text{Aut}(G)$;*
- (b) *G is of the form (euc1), or (euc2), or (hyp1'), or (hyp2'), or (hyp3').*

The complete characterization of compression for mixed type of surface groups remains still an open problem, which seems somehow trickier.

Proof of Theorem 1.13. (a) \Rightarrow (b) is trivial.

For (b) \Rightarrow (c), we apply Theorem 1.11 and remove the case (euc3) by using the counter-example given in Proposition 3.1 from Section 3.

Finally, to see that (c) \Rightarrow (a), we first note that, by elementary results about abelian groups, every subgroup of G is inert in G (and hence compressed in G) in the case (euc1); the same is true for the case (euc2), as proved in Proposition 2.2 from Section 2; finally, for the case (euc4), in Section 4 we use a property of fixed points of endomorphisms to show that $\text{Fix } \phi$ is compressed in G for any $\phi \in \text{End}(G)$; see Corollary 4.3. \square

Proof of Theorem 1.14. (a) \Rightarrow (b) is trivial.

For (b) \Rightarrow (c), we apply Theorem 1.13 and remove the case (euc4) by using the counter-example given in Proposition 4.4 from Section 4.

Finally, in abelian groups every subgroup is inert, and the same is true for the case (euc2), as proved in Proposition 2.2; this proves (c) \Rightarrow (a). \square

2. THE CASE (EUC2)

The goal of this section is to show that every subgroup of

$$G = NS_2 \times \mathbb{Z}_2^q = \langle a, b \mid bab^{-1}a \rangle \times \prod_{j=1}^q \langle d_j \mid d_j^2 \rangle,$$

where $q \geq 0$, is inert (and hence compressed) in G . We need the following lemma, which follows from standard arguments.

Lemma 2.1. *Let A, B be two groups, let $G = A \times B$, let $\pi: G \rightarrow A$ be the natural projection (with $\ker \pi = B$), and let $H \leq G$ be a subgroup such that $H \cap B \leq Z(H)$. If $\pi|_H: H \rightarrow \pi H \leq A$ splits then $H \simeq \pi H \times (H \cap B)$.*

Proof. Let $H \leftarrow \pi H: \alpha$ be a morphism such that $\pi\alpha = \text{Id}_{\pi H}$, and consider

$$\begin{aligned} \Psi: H &\rightarrow \pi H \times (H \cap B) \\ h &\mapsto \left(\pi h, (\alpha\pi h)^{-1} \cdot h \right). \end{aligned}$$

It is well-defined since $\pi((\alpha\pi h)^{-1} \cdot h) = (\pi\alpha\pi h)^{-1} \cdot (\pi h) = 1$ and so, $(\alpha\pi h)^{-1} \cdot h \in \ker \pi|_H = H \cap B$. It is a morphism since

$$\begin{aligned} \Psi(h_1 h_2) &= \left(\pi(h_1 h_2), (\alpha\pi(h_1 h_2))^{-1} \cdot (h_1 h_2) \right) \\ &= \left((\pi h_1) \cdot (\pi h_2), (\alpha\pi h_2)^{-1} \cdot (\alpha\pi h_1)^{-1} \cdot h_1 \cdot h_2 \right) \\ &= \left((\pi h_1) \cdot (\pi h_2), (\alpha\pi h_1)^{-1} \cdot h_1 \cdot (\alpha\pi h_2)^{-1} \cdot h_2 \right) \\ &= \Psi h_1 \cdot \Psi h_2, \end{aligned}$$

where the middle step is valid because $(\alpha\pi h_1)^{-1} \cdot h_1 \in H \cap B \leq Z(H)$, by hypothesis. Finally, it is straightforward to see that $\pi H \times (H \cap B) \rightarrow H$, $(x, y) \mapsto (\alpha x) \cdot y$ is a homomorphism and it is the inverse of Ψ . Hence, Ψ is bijective. \square

Proposition 2.2. *Every subgroup $H \leq G = NS_2 \times \mathbb{Z}_2^q$, $q \geq 0$, is inert (and hence compressed) in G .*

Proof. First, note that any subgroup of NS_2 is isomorphic to either $1, \mathbb{Z}, \mathbb{Z}^2$, or NS_2 ; it is then immediate that any subgroup of NS_2 is inert in NS_2 ; this is the case $q = 0$.

Suppose $q \geq 1$, fix an arbitrary $H \leq G = NS_2 \times \mathbb{Z}_2^q$, and let us see it satisfies the conditions of Lemma 2.1. The fact $H \cap \mathbb{Z}_2^q \leq Z(H)$ is obvious because \mathbb{Z}_2^q is abelian. To see that the restricted short exact sequence $1 \rightarrow H \cap \mathbb{Z}_2^q \rightarrow H \rightarrow \pi H \rightarrow 1$ splits, let us distinguish the isomorphism type of $\pi H \leq NS_2$: if $\pi H \simeq 1, \mathbb{Z}$ then it is free and the sequence clearly splits; if $\pi H \simeq \mathbb{Z}^2, NS_2$, take two elements $u, v \in \pi H$ which present πH , say $\pi H \simeq \langle u, v \mid uv = vu \rangle$ or $\pi H \simeq \langle u, v \mid vuv^{-1}u \rangle$, respectively, and choose arbitrary preimages $ue_u, ve_v \in H$, where $e_u, e_v \in \mathbb{Z}_2^q$ and so satisfy $e_u^2 = e_v^2 = 1$. We claim that, in both cases, $\alpha: \pi H \rightarrow H$, $u \mapsto ue_u, v \mapsto ve_v$ splits the sequence. In fact, it is a well-defined morphism because, if $uv = vu$ in πH then certainly $ue_u \cdot ve_v = ve_v \cdot ue_u$ in G ; and if $vuv^{-1}u = 1$ in πH then certainly $ve_v \cdot ue_u \cdot (ve_v)^{-1} \cdot ue_u = vuv^{-1}u \cdot e_u^2 = 1$ in G . Hence, by Lemma 2.1, $H \simeq \pi H \times (H \cap \mathbb{Z}_2^q)$.

Now take an arbitrary $R \leq G$. By the previous paragraph, we also have $R \simeq \pi R \times (R \cap \mathbb{Z}_2^q)$ and $H \cap R \simeq \pi(H \cap R) \times (H \cap R \cap \mathbb{Z}_2^q)$. Then,

$$\text{rk}(H \cap R) = \text{rk}(\pi(H \cap R)) + \text{rk}(H \cap R \cap \mathbb{Z}_2^q) \leq \text{rk}(\pi R) + \text{rk}(R \cap \mathbb{Z}_2^q) = \text{rk}(R),$$

where the inequality $\text{rk}(\pi(H \cap R)) \leq \text{rk}(\pi R)$ is true because $\pi(H \cap R) \leq \pi R \leq NS_2$, and subgroups of NS_2 are compressed in NS_2 . \square

Remark 2.3. We want to direct reader's attention to the fact that elements in \mathbb{Z}_2^q all have order 2; this played a crucial role in the above argument because, whatever preimage $ue_u \in H$ we choose for $u \in \pi H$ satisfies $e_u^2 = 1$, and this was necessary to make α well-defined in the case $\pi H \simeq NS_2$. This small detail is important in light of the following example.

We observe that the result stated in Proposition 2.2 is *not* true if we remove this order two relation: in general, subgroups of $G = NS_2 \times \mathbb{Z} = \langle a, b \mid bab^{-1}a \rangle \times \langle c \rangle$ are *no longer* inert, neither even compressed in G . Consider $R = \langle ac, b \rangle \leq G$; it contains $a^{-1}c = b(ac)b^{-1}$ and so also contains $a^2 = (ac)(a^{-1}c)^{-1}$ and $c^2 = (ac)(a^{-1}c)$; therefore, $H = \langle a^2, b^2, c^2 \rangle = \langle a^2, b^2 \rangle \times \langle c^2 \rangle \simeq \mathbb{Z}^3$ is contained in $R \leq G$ and so, it is not compressed in G . (Note that, inserting this R into the argument in Proposition 2.2, the restriction of the projection $\pi: G \rightarrow NS_2$ to R , say $\pi|_R: R \rightarrow \pi R = NS_2$, $ac \mapsto a, b \mapsto b$ *does not* split: all elements $r \in R$ satisfy $|r|_c = |r|_a$ (which only makes sense modulo 2) and so all preimages of a in R have odd, and so non-trivial, c -exponent; hence, the relation $bab^{-1}a$ cannot be preserved up in R .)

This remark tells us that a general approach like we did in the case (euc2) will not work in the case (euc4). So, in Section 4, we will be forced to use specific properties of fixed subgroups of endomorphisms.

3. THE CASE (EUC3)

In this section, we consider the case (euc3), $G = NS_2 \times \mathbb{Z}^p \times \mathbb{Z}_2$ for some $p \geq 1$, and exhibit an automorphism $\phi \in \text{Aut}(G)$ such that $\text{Fix } \phi$ is not compressed (and hence not inert) in G .

Put $G = NS_2 \times \mathbb{Z}^p \times \mathbb{Z}_2 = \langle a, b \mid bab^{-1}a \rangle \times \prod_{i=1}^p \langle c_i \rangle \times \langle d \mid d^2 \rangle$. Since $ba^n = a^{-n}b$ for $n \in \mathbb{Z}$, and $c_i, d \in Z(G)$ for $i = 1, \dots, p$, we see that each $g \in G$ can be written as $g = a^{k_0} b^{k_1} c_1^{n_1} \cdots c_p^{n_p} d^{k_2}$, for some unique $k_0, k_1, n_1, \dots, n_p \in \mathbb{Z}$ and unique $k_2 \in \mathbb{Z}_2$.

Consider $\phi: G \rightarrow G$ given by $a \mapsto ad$, $b \mapsto ba$, $c_1 \mapsto c_1 d$, $c_i \mapsto c_i^{-1}$ ($i = 2, \dots, p$), $d \mapsto d$.

Proposition 3.1. *ϕ is a well-defined automorphism of $G = NS_2 \times \mathbb{Z}^p \times \mathbb{Z}_2$, $p \geq 1$, and $\text{Fix } \phi$ is not compressed in G .*

Proof. It is straightforward to see that the defining relations from G are preserved and so, ϕ is a well-defined endomorphism of G . To see it is an automorphism, one just checks that $\psi: G \rightarrow G$, $a \mapsto ad$, $b \mapsto ba^{-1}d$, $c_1 \mapsto c_1 d$, $c_i \mapsto c_i^{-1}$ ($i = 2, \dots, p$), $d \mapsto d$ is again well defined and satisfies $\phi\psi = \psi\phi = \text{Id}$.

Now let us prove that $\text{Fix } \phi = \langle a^2, b^2, ac_1, d \rangle$. The inclusion \supseteq is straightforward. Let $g = a^{k_0} b^{k_1} c_1^{n_1} c_2^{n_2} \cdots c_p^{n_p} d^{k_2} \in G$ be fixed by ϕ ; we have

$$a^{k_0} b^{k_1} c_1^{n_1} c_2^{n_2} \cdots c_p^{n_p} d^{k_2} = g = \phi g = a^{k_0 - \delta(k_1)} b^{k_1} c_1^{n_1} c_2^{-n_2} \cdots c_p^{-n_p} d^{k_0 + n_1 + k_2},$$

where $\delta(k)$ is 1 if k is odd, and 0 if k is even. So, $n_2 = \cdots = n_p = 0$, and both k_1 and $k_0 + n_1$ must be even, therefore $k_0 - n_1$ is also even and $g = a^{k_0} b^{k_1} c_1^{n_1} c_2^{n_2} \cdots c_p^{n_p} d^{k_2} = a^{k_0 - n_1} b^{k_1} (ac_1)^{n_1} d^{k_2} \in \langle a^2, b^2, ac_1, d \rangle$ (recall that b^2 commutes with a).

Finally, note that $\text{Fix } \phi = \langle a^2, b^2, ac_1 \rangle \times \langle d \rangle \cong \mathbb{Z}^3 \times \mathbb{Z}_2$ has rank 4 but, $\text{Fix } \phi \leq \langle ac_1, b, d \rangle$ since $a^2 = (ac_1)b(ac_1)^{-1}b^{-1}$. Hence, $\text{Fix } \phi$ is not compressed in G . \square

4. THE CASE (EUC4)

In this section, we concentrate on case (euc4), i.e.,

$$G = NS_2^\ell \times \mathbb{Z}^p = \prod_{i=1}^{\ell} \langle a_i, b_i \mid b_i a_i b_i^{-1} a_i \rangle \times \prod_{j=1}^p \langle c_j \rangle,$$

$\ell \geq 1$, $p \geq 0$. We shall prove that the fixed subgroup of any endomorphism of G is compressed in G (despite the fact that, except for the case $\ell = 1$, $p = 0$, G contains some subgroups which are not compressed; see Remark 2.3). When $\ell + p \geq 2$, we shall also exhibit an automorphism $\phi \in \text{Aut}(G)$ such that $\text{Fix } \phi$ is not inert in G .

With this notation, consider $G \supseteq N = \langle a_1, \dots, a_\ell \rangle \cong \mathbb{Z}^\ell$, and the commutator subgroup, $G \supseteq G' = \langle a_1^2, \dots, a_\ell^2 \rangle \cong \mathbb{Z}^\ell$. Observe that every $g \in G'$ has a unique square root in G , which happens to belong to N , and we denote it by $\sqrt{g} \in N$: in fact, in each NS_2 direct factor of G , the square of an element with non-zero total number of b_i 's has non-zero total number of b_i 's. (In general, however, roots are not unique in NS_2 , since $(a_i^r b_i)^2 = b_i^2$ for every $r \in \mathbb{Z}$.)

Lemma 4.1. *Let $\varphi: G_1 \rightarrow G_2$ be a group morphism. If a subgroup $H \leq G_1$ satisfies $\text{rk}(\varphi(H)) = \text{rk}(H)$, and $\varphi(H)$ is compressed in $\text{Im}(\varphi)$ then, H is compressed in G_1 .*

Proof. For every $H \leq K \leq G_1$, we have $\varphi(H) \leq \varphi(K) \leq \text{Im}(\varphi) \leq G_2$; therefore, $\text{rk}(H) = \text{rk}(\varphi(H)) \leq \text{rk}(\varphi(K)) \leq \text{rk}(K)$. \square

Proposition 4.2. *Let $G = NS_2^\ell \times \mathbb{Z}^p$, $\ell \geq 1$, $p \geq 0$. If a subgroup $H \leq G$ satisfies that $\sqrt{h} \in H$ for every $h \in H \cap G'$, then H is compressed in G .*

Proof. Let $\{h_1, \dots, h_k\}$ be a free-abelian basis of $H \cap G' \simeq \mathbb{Z}^k$, $k \leq \ell$. From the hypothesis on H , it follows that $\{\sqrt{h_1}, \dots, \sqrt{h_k}\}$ is a free-abelian basis of $H \cap N \simeq \mathbb{Z}^k$. Therefore, $(H \cap N)/(H \cap G') \simeq \mathbb{Z}_2^k$.

On the other hand $H/(H \cap N) \leq G/N \simeq \mathbb{Z}^{\ell+p}$, so $H/(H \cap N) \simeq \mathbb{Z}^r$ for some $r \leq \ell + p$; also, $H/(H \cap G') \leq G/G' \simeq \mathbb{Z}_2^\ell \times \mathbb{Z}^{\ell+p}$, so $H/(H \cap G') \simeq \mathbb{Z}_2^s \times \mathbb{Z}^t$ for some $s \leq \ell$ and $t \leq \ell + p$. But

$$\mathbb{Z}^r \simeq H/(H \cap N) \simeq \frac{H/(H \cap G')}{(H \cap N)/(H \cap G')} \simeq \frac{\mathbb{Z}_2^s \times \mathbb{Z}^t}{\mathbb{Z}_2^k}$$

so, $s = k$ and $t = r$. Then, for the abelianization map $\rho: G \rightarrow G/G'$, we have $\text{rk}(\rho(H)) = \text{rk}(H/(H \cap G')) = s + t = k + r = \text{rk}(H \cap N) + \text{rk}(H/(H \cap N)) \geq \text{rk}(H)$ and hence, $\text{rk}(\rho(H)) = \text{rk}(H)$. By Lemma 4.1, H is compressed in G . \square

Corollary 4.3. *Let $G = NS_2^\ell \times \mathbb{Z}^p$, $\ell \geq 1$, $p \geq 0$. For every $\phi \in \text{End}(G)$, $\text{Fix } \phi$ is compressed in G .*

Proof. By Proposition 4.2, it suffices to show that $\sqrt{h} \in \text{Fix } \phi$, for every $h \in \text{Fix } \phi \cap G'$. But this follows from the above observation that \sqrt{h} is the unique element in G with square equal to h (hence, if h is fixed by ϕ , \sqrt{h} must also be fixed by ϕ). \square

Finally, in this case (euc4), we give examples of automorphisms $\phi \in \text{Aut}(G)$ such that $\text{Fix } \phi$ is not inert in G (except, of course, for the case $\ell = 1$, $p = 0$, which is included in (euc2)).

Proposition 4.4. *Let $G = NS_2^\ell \times \mathbb{Z}^p$, $\ell \geq 1$, $\ell + p \geq 2$. There exists an automorphism $\phi \in \text{Aut}(G)$ such that $\text{Fix } \phi$ is not inert in G .*

Proof. The condition $\ell + p \geq 2$ (necessary in order to avoid the case $G = NS_2$, for which we already know that every subgroup is inert) is equivalent to $p \geq 1$ or $\ell \geq 2$, just meaning that we have at least two direct factors in G .

Suppose $\ell = p = 1$, i.e., $G = NS_2 \times \mathbb{Z} = \langle a, b \mid bab^{-1}a \rangle \times \langle c \rangle$. Consider $\phi_1: G \rightarrow G$, $a \mapsto a$, $b \mapsto ba$, $c \mapsto c$; straightforward calculations show that it is a well-defined automorphism $\phi_1 \in \text{Aut}(G)$, and that $\text{Fix } \phi_1 = \langle a, b^2, c \rangle \cong \mathbb{Z}^3$. But this is not inert in G because $\text{Fix } \phi_1 \cap \langle ac, b \rangle = \langle ac, a^2, b^2 \rangle \cong \mathbb{Z}^3$ (the inclusion \geq follows from direct inspection, and \leq is a consequence of $\langle ac, a^2, b^2 \rangle$ being an index 2 subgroup of $\langle a, b^2, c \rangle$).

Now, suppose $\ell = 2$, $p = 0$, i.e., $G = (NS_2)^2 = \langle a_1, b_1 \mid b_1 a_1 b_1^{-1} a_1 \rangle \times \langle a_2, b_2 \mid b_2 a_2 b_2^{-1} a_2 \rangle$. Consider $\phi_2: G \rightarrow G$, $a_1 \mapsto a_1$, $b_1 \mapsto b_1 a_1$, $a_2 \mapsto a_2$, $b_2 \mapsto b_2^{-1}$; again, straightforward calculations show that it is a well-defined automorphism $\phi_2 \in \text{Aut}(G)$, and that $\text{Fix } \phi_2 = \langle a_1, b_1^2, a_2 \rangle \cong \mathbb{Z}^3$. By the same argument as in the previous paragraph, this is not inert in G .

The general case follows easily by taking the identity in all the following direct factors beyond the first two. \square

Remark 4.5. The concepts of compression and inertia are quite delicate and sensible to the ambient group. In the last sentence of the previous proof, we used the straightforward fact that, for $\phi \in \text{Aut}(G_1)$, if $\text{Fix}(\phi)$ is not inert in G_1 then $\text{Fix}(\phi \times id)$ is not inert in $G_1 \times G_2$. However, the converse is not true: we proved that $\langle a, b^2 \rangle \times \langle c \rangle$ is *not* inert in $NS_2 \times \mathbb{Z}$, while $\langle a, b^2 \rangle$ is inert in NS_2 , by Proposition 2.2.

Acknowledgements. The first named author is partially supported by the National Natural Science Foundation of China (No. 11571246); the second by the Spanish Agencia Estatal de Investigación, through grant MTM2017-82740-P (AEI/FEDER, UE), and also by the “María de Maeztu” Programme for Units of Excellence in R&D (MDM-2014-0445); finally, the third named author is partially supported by the National Natural Science Foundation of China (No. 11771345).

REFERENCES

- [1] G. Bergman, “Supports of derivations, free factorizations and ranks of fixed subgroups in free groups”, *Trans. Amer. Math. Soc.* **351** (1999), 1531–1550.
- [2] M. Bestvina and M. Handel, “Train tracks and automorphisms of free groups”, *Annals of Mathematics* **135** (1992), 1–51.
- [3] W. Dicks and E. Ventura, “The group fixed by a family of injective endomorphisms of a free group”, *Contemporary Mathematics* **195**, AMS, Providence (1996).
- [4] W. Jaco and P. Shalen, “Surface homeomorphisms and periodicity”, *Topology*, **16** (1977), 347–367.
- [5] B. Jiang, S. Wang, and Q. Zhang, “Bounds for fixed points and fixed subgroups on surfaces and graphs”, *Alg. Geom. Topology* **11** (2011), 2297–2318.
- [6] J. Lin and S. Wang, “Fixed subgroups of automorphisms of hyperbolic 3-manifold groups”, *Topology Appl.* **173** (2014), 175–187.
- [7] A. Martino and E. Ventura, “Fixed subgroups are compressed in free groups”, *Communications in Algebra* **32** (2004), no. 10, 3921–3935.
- [8] J. Nielsen, “Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen”, *Acta Math.* **50** (1927), 189–358.
- [9] J. Nielsen, “Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen. II”, *Acta Math.* **53** (1929), 1–76.
- [10] J. Wu and Q. Zhang, “The group fixed by a family of endomorphisms of a surface group”, *J. Algebra* **417** (2014), 412–432.
- [11] Q. Zhang, “Bounds for fixed points on Seifert manifolds”, *Topology and its Applications* **159** (2012), 3263–3273.
- [12] Q. Zhang, “The fixed subgroups of homeomorphisms of Seifert manifolds”, *Acta Math. Sin. (Engl. Ser.)* **31** (2015), 797–810.
- [13] Q. Zhang, E. Ventura, and J. Wu “Fixed subgroups are compressed in surface groups”, *Internat. J. Algebra Comput.* **25** (2015), 865–887.
- [14] H. Zieschang, “Über einfache Kurven auf Vollbrezeln”, *Abh. Math. Sem. Univ. Hamburg* **25** (1961/1962), 231–250.

DEPARTMENT OF MATHEMATICS, SOOCHOW UNIVERSITY, SUZHOU 215006, CHINA

Email address: wujianchun@suda.edu.cn

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT POLITÈCNICA DE CATALUNYA, CATALONIA

Email address: enric.ventura@upc.edu

SCHOOL OF MATHEMATICS AND STATISTICS, XI’AN JIAOTONG UNIVERSITY, XI’AN 710049, CHINA

Email address: zhangq.math@mail.xjtu.edu.cn